Andean Geology 53 (3): 810-813. September, 2026
doi: 10.5027/andgeoV53n3-3959

Andean Geology
www.andeangeology.cl

COMMENT

Comment on "Applied geophysics to morphostructurally characterize a hydrothermal system: La Laja, San Juan (Argentina)" by Sottile *et al*. (2026), Andean Geology 53 (1): 113-135

Ricardo A. Astini[1, 2]

[1] *Facultad de Ciencias Exactas Físicas y Naturales, Universidad Nacional de Córdoba, av. Vélez Sarsfield 1611, Córdoba, X5016GCA, Argentina.*
ricardo.astini@unc.edu.ar

[2] *Centro de Investigaciones en Ciencias de la Tierra, CONICET, Ingeniero Ismael Bordabehere, av. Haya de la Torre, Córdoba, X5016GCA, Argentina.*

The contribution by Sottile *et al*. (2026) represents a valuable addition to the understanding of the La Laja system (San Juan, Argentina), particularly through the integration of geophysical, structural and geomorphological data. The identification of structurally controlled permeability and reservoir architecture provides an important framework for interpreting fluid circulation.

In this context, it may be useful to further refine the classification of the La Laja system. While the authors refer to it as a hydrothermal system, the available evidence appears to be more consistent with a low-temperature geothermal system.

This distinction is not merely semantic, as it involves differences in heat sources, fluid origin, and hydrological functioning. The term "hydrothermal system" encompasses a broad range of geothermal settings, including both magmatically driven systems and those controlled by regional heat flow and hydrogeological circulation (Hochstein, 1990; Dickson and Fanelli, 2004; Sowiżdżał, 2022). In contrast, low-temperature geothermal systems are typically characterized by conductive or conduction-dominated heat transfer, meteoric-dominated recharge, and structurally controlled flow paths, commonly linked to regional groundwater circulation systems and relatively low enthalpy conditions (Ingebritsen *et al*., 2006; Moeck, 2014; Jalilinasrabady *et al*., 2022).

The geophysical and morphostructural results presented by Sottile *et al.* (2026), particularly the role of active faults in controlling permeability and the apparent lack of association with shallow magmatic bodies, are consistent with this interpretation. This view is also in agreement with previous studies (*e.g*., Orozco Chirino *et al*., 2021), that interpret La Laja as a system where meteoric waters infiltrate at higher elevations, circulate at depth, and are heated under regional geothermal gradients before ascending along structural pathways through the largely Paleozoic and Cenozoic sedimentary pile of the Precordillera, including a thick section of Cambrian and Ordovician carbonates (Astini *et al*., 2005). The regional geodynamic context further supports this interpretation. The La Laja system is located within the Pampean flat-slab segment, where arc magmatism was largely suppressed since the Miocene, and any residual magmatic activity was spatially displaced to the east (*e.g*., Kay and Mpodozis, 2002; Ramos *et al*., 2002).

Clarifying this distinction is particularly relevant when considering travertine-forming systems, which can be interpreted in terms of two end-member

models: volcanic-hydrothermal and artesian-karstic. However, these systems are better understood as the result of three partially decoupled controls: (1) heat source (magmatic vs. geothermal), (2) CO_2 source (mantle, crustal, or biogenic; *e.g*., Kerrick, 2001), and (3) hydrological regime (meteoric vs. deep circulation).

Recognizing La Laja as a low-temperature geothermal system has important implications. It supports a model dominated by regional hydrology and structural control, rather than magmatic input, and provides a more consistent framework for interpreting fluid sources, CO_2 origin, and travertine deposition. In this context, travertine distribution and dispersal patterns are expected to be primarily controlled by structural pathways and basin-scale hydrology, which, in turn, influence the spatial organization of facies and the resulting heterogeneity of carbonate accumulations (*e.g*., Pentecost, 2005; Gandin and Capezzuoli, 2014; Mors *et al*., 2019; Astini *et al*., 2025).

In addition, the travertine system at La Laja displays a wide range of mound morphologies (Astini *et al*., 2016; also see figures 5, 7, 9 and 10 in Sottile *et al*., 2026), which may provide further insights into the underlying hydrodynamic and structural controls. Field observations indicate the coexistence of conical subaerial mounds with high aspect ratios and relatively stable central vent positions, as well as broader, low-relief, discoidal bodies with depressed central areas that likely functioned as shallow pools with subaqueous precipitation.

Such morphological variability has been shown to reflect differences in discharge rate, fluid pressure, vent stability, and self-sealing processes driven by carbonate precipitation (*e.g*., Luo *et al.*, 2022). In this context, the integration of geophysical data with detailed morphological and sedimentological observations could help constrain the controlling parameters of mound growth, including the role of structurally focused flow, temporal variability in discharge, and feedback mechanisms between permeability and precipitation. This approach may provide a valuable pathway to link subsurface structure with surface expression in geothermal travertine systems. This has direct implications not only for the understanding of depositional architecture, but also for the evaluation of travertine bodies as industrial carbonate resources (dimension stone and related applications), where facies distribution, porosity, and diagenetic overprint determine their quality and exploitability.

On a broader scale, the same controls govern the spatial organization of facies and the resulting heterogeneity of carbonate accumulations, which are widely used as analogues in reservoir characterization (*e.g*., Claes *et al*., 2015; Della Porta, 2015; Ronchi and Cruciani, 2015; De Boever *et al*., 2016). In these systems, permeability architecture reflects the interplay between fluid flow, degassing, and accommodation space, leading to strong facies-controlled heterogeneity and multi-scale variability in petrophysical properties (Lucia, 1999; Mancini *et al*., 2019).

This perspective also provides a useful framework to further explore the architecture of the feeding system (plumbing system) and its role in controlling the spatial distribution and dispersal of surface depositional products. In particular, it enables a more robust genetic interpretation of the facies mosaic, linking fluid pathways, degassing processes, and accommodation space to structurally controlled discharge zones. These aspects appear to be closely tied to geophysical signatures associated with active fault systems, highlighting the broader implications of the results presented by Sottile *et al*. (2026). It also has practical implications for resource assessment and hazard evaluation, as low-temperature systems differ significantly from magmatically driven hydrothermal systems in both geothermal potential and associated risks (Dickson and Fanelli, 2004). In contrast, magmatic-hydrothermal systems are more commonly associated with metal transport and deposition due to higher temperatures and more intense fluid-rock interaction, and therefore have a greater potential for ore-forming processes (*e.g*., Robb, 2005; Simmons *et al*., 2005). Recognizing the geothermal (non-magmatic) nature of La Laja thus constrains both its economic potential and its genetic interpretation.

Building on the valuable geophysical and morphostructural framework provided by Sottile *et al*. (2026), further integration with complementary datasets could help refine the conceptual model of the La Laja system. In particular, the incorporation of hydrogeochemical and isotopic tracers (*e.g*., $\delta^{13}C$-CO_2, $\delta^{18}O$-H_2O) would allow better constraining fluid sources, residence times, and fluid-rock interaction processes (*e.g*., Kerrick, 2001; Ingebritsen *et al*., 2006). Likewise, detailed sedimentological and facies analysis of associated travertine deposits currently

under study could provide independent insights into fluid pathways, degassing dynamics, and the spatial organization of depositional systems.

The present perspective highlights the importance of integrating geothermal systems within broader basin-scale hydrological models, particularly in arid regions such as San Juan, where groundwater circulation and structural controls play a dominant role. Integrating the structural framework with basin-scale hydrogeological models would contribute to a more comprehensive understanding of recharge areas, circulation depths, and discharge patterns, particularly in the context of the arid Andes. Such an approach may help bridge geophysical observations with surface expressions and improve the predictive understanding of similar systems in the region.

References

Astini, R.A.; Dávila, F.M.; López Gamundí, O.; Gómez, F.; Collo, G.; Ezpeleta, M.; Martina, F.; Ortiz, A. 2005. Cuencas de la región precordillerana. *In* Frontera Exploratoria de la Argentina (Chebli, G.; Spalletti, L.; editors). Instituto Argentino del Petróleo y del Gas: 115-145. Buenos Aires.

Astini, R.; Gómez, F.; De Maestri, E. 2016. Los travertinos cuaternarios de La Laja (San Juan): morfología, estratigrafía, facies y paleoambientes. *In* Congreso Latinoamericano de Sedimentología, No. 7, Reunión Argentina de Sedimentología, No. 15, Libro de resúmenes: p. 28. La Pampa.

Astini, R.A.; Mors, R.A.; Sariago, R.A.; Druetta, H.S. 2025. Carbonatos asociados con fenómenos hidrotermales y póstumos al volcanismo en la región del cerro Tiporco. *In* Geología y Recursos Naturales de la provincia de San Luis (Costa, C.H.; editor). Congreso Geológico Argentino, No. 22, Relatorio: 451-478, San Luis.

Claes, H.; Soete, J.; Van Noten, K.; El Desouky, H.; Marques Erthal, M.; Vanhaecke, F.; Özkul, M.; Swennen, R. 2015. Sedimentology, three-dimensional geobody reconstruction and carbon dioxide origin of Pleistocene travertine deposits in the Ballık area (south-west Turkey). Sedimentology 62 (5): 1408-1445. https://doi.org/10.1111/sed.12188

De Boever, E.; Foubert, A.; Oligschlaeger, D.; Claes, S.; Soete, J.; Bertier, P.; Özkul, M.; Virgone, A.; Swennen, R. 2016. Multiscale approach to (micro) porosity quantification in continental spring carbonate facies: Case study from the Cakmak quarry (Denizli, Turkey). Geochemistry, Geophysics, Geosystems 17 (7): 2922-2939.

Della Porta, G. 2015. Carbonate build-ups in lacustrine, hydrothermal and fluvial settings: comparing depositional geometry, fabric types and geochemical signature. *In* Microbial Carbonates in Space and Time: Implications for Global Exploration and Production (Bosence, D.W.J.; Gibbons, K.A.; Le Heron, D.P.; Morgan, W.A.; Pritchard, T.; Vining, B.A.; editors). Geological Society, Special Publications 418: 17-68. London.

Dickson, M.H.; Fanelli, M. 2004. What is Geothermal Energy? International Geothermal Association (IGA): 15 p. Pisa.

Gandin, A.; Capezzuoli, E. 2014. Travertine: distinctive depositional fabrics of carbonates from thermal spring systems. Sedimentology 61 (1): 264-290. https://doi.org/10.1111/sed.12087

Hochstein, M.P. 1990. Classification and assessment of geothermal resources. *In* Small Geothermal Resources: A Guide to Development and Utilization (Dickson, M.H.; Fanelli, M.; editors). UNITAR/UNDP Centre on Small Energy Sources: 31-59. Roma.

Ingebritsen, S.E.; Sanford, W.E.; Neuzil, C.E. 2006. Groundwater in Geologic Processes (2nd edition). Cambridge University Press: 536 p. Cambridge.

Jalilinasrabady, S.; Itoi, R.; Setyawan, A. 2022. Geothermal Resources: Briefing on Geothermal Systems and Their Classification. *In* Geothermal Energy: An Overview on Resources and Potential (Setyawan, A.; Setyawan; A. Jalilinasrabady, S.; Pambudi, N.A.; Itoi, R.; editors). Springer: 1-25. Cham.

Kay, S.M.; Mpodozis, C. 2002. Magmatism as a probe to the Neogene shallowing of the Nazca plate beneath the modern Chilean flat-slab. Journal of South American Earth Sciences 15 (1): 39-57.

Kerrick, D.M. 2001. Present and past nonanthropogenic CO_2 degassing from the solid earth. Reviews of Geophysics 39 (4): 565-585.

Lucia, F.J. 1999. Carbonate Reservoir Characterization. Springer-Verlag: 226 p. Berlin.

Luo, L.; Wen, H.; Brogi, A.; Capezzuoli, E. 2022. Factors controlling the geometry of travertine mounds: Insights from Heinitang (China). Sedimentology 69 (4): 1519-1546.

Mancini, A.; Capezzuoli, E.; Erthal, M.; Swennen, R. 2019. Hierarchical approach to define travertine depositional systems: 3D conceptual morphological model and possible applications. Marine and Petroleum Geology 103: 549-563. https://doi.org/10.1016/j.marpetgeo.2019.02.021

Moeck, I.S. 2014. Catalog of geothermal play types based on geologic controls. Renewable and Sustainable Energy Reviews 37: 867-882.

Mors, R.A.; Astini, R.A.; Gómez, F.J. 2019. Coexisting active travertines and tufas in the southeastern border of the Puna plateau. Sedimentary Geology 389: 200-217.

Orozco Chirino, P.G.; Astini, R.A.; Presa, J.M.; Alvarado, P.M.; Venerdini, A.L.; Rivas Bilbao, A.C.; Pérez Luján, S.B.; Ortiz, G.F.; Peralta Arnold, Y.J; Sáez, M. 2021. The La Laja spring system in the Argentine Precordillera: A conceptual model based on geochemical and isotopic data. Journal of South American Earth Sciences 109:103241. https://dx.doi.org/10.1016/j.jsames.2021.103241

Pentecost, A. 2005. Travertine. Springer: 445 p. Berlin.

Ramos, V.A.; Cristallini, E.O.; Pérez, D.J. 2002. The Pampean flat-slab of the Central Andes. Journal of South American Earth Sciences 15 (1): 59-78.

Robb, L. 2005. Introduction to Ore-Forming Processes; Blackwell: 373 p. Malden, MA.

Ronchi, P.; Cruciani, F. 2015. Continental carbonates as a hydrocarbon reservoir, an analog case study from the travertine of Saturnia, Italy. American Association Petroleum of Geologists Bulletin 99 (4): 711-734.

Simmons, S.F.; White, N.C.; John, D.A. 2005. Geological characteristics of epithermal precious and base metal deposits. *In* Economic Geology One Hundredth Anniversary Volume (1905-2005) (Hedenquist, J.W.; Thompson, J.F.H.; Goldfarb, R.J.; Richards, J.P.; editors). Society of Economic Geologists: 485-522. Littleton.

Sottile, A.; Rodríguez, A.; Winocur, D.; Grigolo, M; Martínez, M. 2026. Applied geophysics to morphostructurally characterize a hydrothermal system: La Laja, San Juan (Argentina). Andean Geology 53 (1): 113-135. https://dx.doi.org/10.5027/andgeoV53n1-3784

Sowiżdżał, A. 2022. Geothermal Systems-An Overview. Energies 15: 6377. https://doi.org/10.3390/en15176377

Manuscript received: March 20, 2026; revised/accepted: April 07, 2026; available online: April 07, 2026.